

Group-theoretical construction of extended baryon operators

LHP Collaboration: S. Basak^a, R. Edwards^b, R. Fiebig^c, G.T. Fleming^{db}, U.M. Heller^e, C. Morningstar^{f*}, D. Richards^b, I. Sato^a, S. Wallace^a.

^aDepartment of Physics, University of Maryland, College Park, MD 20742, USA

^bThomas Jefferson National Accelerator Facility, Newport News, VA 23606, USA

^cPhysics Department, Florida International University, Miami, FL 33199, USA

^dSloane Physics Laboratory, Yale University, New Haven, CT 06520, USA

^eAmerican Physical Society, Ridge, NY 11961-9000, USA

^fDepartment of Physics, Carnegie Mellon University, Pittsburgh, PA 15213, USA

The design and implementation of large sets of spatially extended baryon operators for use in lattice simulations are described. The operators are constructed to maximize overlaps with the low-lying states of interest, while minimizing the number of sources needed in computing the required quark propagators.

1. OVERVIEW

The Lattice Hadron Physics Collaboration (LHPC) was given a charge a few years ago by the late Nathan Isgur to extract as much as possible of the low-lying spectrum of baryon resonances from Monte Carlo simulations in lattice QCD. To accomplish this ambitious task, we have been pursuing two different approaches to constructing baryon interpolating field operators. Progress with the first of these approaches is reported elsewhere in this conference[1,2]. In this talk, developments in the design and implementation of extended baryon operators using the second approach are outlined. Initial efforts using this second method were described at the Tsukuba symposium last year[3]. Some details presented in the earlier reference will not be repeated in this brief report.

The usual operator construction which mimics the approach one would take in continuous space-time (such as in the QCD sum-rule method) is very cumbersome, especially when tackling an entire spectrum, as opposed to a single state or a few states. Our approach described here is in-

stead to directly combine the physical characteristics of baryons with the symmetries of the lattice regularization of QCD used in simulations. For baryons at rest, our operators are formed using group-theoretical projections onto the irreducible representations (irreps) of the O_h symmetry group of a three-dimensional cubic lattice. There are four two-dimensional irreps $G_{1g}, G_{1u}, G_{2g}, G_{2u}$ and two four-dimensional representations H_g and H_u . The continuum-limit spins J of our states must be deduced by examining degeneracy patterns across the different O_h irreps[3].

Baryons are expected to be rather large objects, and hence, local operators will not suffice. Our approach to constructing extended operators is to use covariant displacements of the quark fields. Displacements in different directions are used to build up the appropriate orbital structure, and displacements of different lengths can build up the needed radial structure. All of our three-quark baryon operators are linear superpositions of gauge-invariant terms of the form

$$\Phi_{\alpha i; \beta j; \gamma k}^{ABC} = \varepsilon_{abc} (\tilde{D}_i^{(p)} \tilde{\psi})_{a\alpha}^A (\tilde{D}_j^{(p)} \tilde{\psi})_{b\beta}^B (\tilde{D}_k^{(p)} \tilde{\psi})_{c\gamma}^C, \quad (1)$$

where A, B, C indicate quark flavor, a, b, c are color indices, α, β, γ are Dirac spin indices, $\tilde{\psi}$ indi-

*Presented by C. Morningstar.

cates a smeared quark field, and $\tilde{D}_j^{(p)}$ denotes the p -link covariant displacement operator in the j -th direction. The three-dimensional gauge-covariant Laplacian is used to smear the quark fields:

$$\begin{aligned}\tilde{\psi}(x) &= (1 + \varrho \tilde{\Delta})^{n_\varrho} \psi(x), \\ \tilde{\Delta}\psi(x) &= \sum_{k=\pm 1, \pm 2, \pm 3} \left(\tilde{U}_k(x) \psi(x + \hat{k}) - \psi(x) \right),\end{aligned}\quad (2)$$

where ϱ and integer n_ϱ are tunable parameters, and $\tilde{U}_\mu(x)$ denotes a link variable smeared using the stout link algorithm[4]. The p -link gauge-covariant displacement operator is defined by

$$\tilde{D}_j^{(p)} O(x) = \tilde{U}_j(x) \dots \tilde{U}_j(x + (p-1)\hat{j}) O(x + p\hat{j}). \quad (4)$$


In the definition of $\Phi_{\alpha i; \beta j; \gamma k}^{ABC}$ above, the displacement directions i, j, k can take values $0, \pm 1, \pm 2, \pm 3$, where the value 0 indicates that no displacement occurs. To simplify matters, the displacement length p is taken to be the same for all displaced quarks in a given term. Ultimately, there are six different spatial orientations that we use, shown in Table 1. The singly-displaced operators are meant to mock up a diquark-quark coupling, and the doubly-displaced and triply-displaced operators are chosen since they favor the Δ -flux and Y -flux configurations, respectively.

Next, we combine the $\Phi_{\alpha i; \beta j; \gamma k}^{ABC}$ into *elemental* operators $B_i^F(t, \mathbf{x})$ having the appropriate flavor structure characterized by isospin, strangeness, *etc.* We work in the $m_u = m_d$ (equal u and d quark masses) approximation, and thus, require that the elemental operators have definite isospin, that is, they satisfy appropriate commutation relations with the isospin operators τ_3, τ_+, τ_- . Since we plan to compute full correlation matrices, we need not be concerned with forming operators according to an $SU(3)$ flavor symmetry. For zero momentum states, we impose translation invariance: $B_i^F(t) = \sum_{\mathbf{x}} B_i^F(t, \mathbf{x})$. Maple code which manipulates Grassmann fields was used to identify maximal sets of linearly independent elemental operators. The numbers of such operators are listed in Table 2.

The final step in our operator construction is to apply group-theoretical projections to obtain

Table 1

Six types of three-quark $\Phi_{\alpha i; \beta j; \gamma k}^{ABC}$ operators used. Smeared quarks fields are shown by solid circles, line segments indicate covariant displacements, and each hollow circle indicates the location of a color ε_{abc} coupling. For simplicity, all displacements have the same length in an operator.

Operator type	Displacement indices
	$i = j = k = 0$
single-site	
	$i = j = 0, k \neq 0$
singly-displaced	
	$i = 0, j = -k, k \neq 0$
doubly-displaced-I	
	$i = 0, j \neq k , jk \neq 0$
doubly-displaced-L	
	$i = -j, j \neq k , jk \neq 0$
triply-displaced-T	
	$ i \neq j \neq k , ijk \neq 0$
triply-displaced-O	

operators which transform irreducibly under all lattice rotation and reflection symmetries:

$$B_i^{\Lambda \lambda F}(t) = \frac{d_\Lambda}{g_{O_h}} \sum_{R \in O_h} D_{\lambda \lambda}^{(\Lambda)}(R)^* U_R B_i^F(t) U_R^\dagger, \quad (5)$$

where Λ refers to an O_h irrep, λ is the irrep row, g_{O_h} is the number of elements in O_h , d_Λ is the dimension of the Λ irrep, $D_{mn}^{(\Lambda)}(R)$ is a Λ representation matrix corresponding to group element R , and U_R is the quantum operator which implements the symmetry operations.

The projections in Eq. (5) are carried out using Maple in the following sequence of steps. (a) A set of M_B linearly-independent elemental operators $B_j^F(t)$ that transform among one another under O_h is identified. (b) We obtain the $M_B \times M_B$ representation matrices $D_{ij}(R)$ which satisfy

$$U_R B_i^F(t) U_R^\dagger = \sum_{j=1}^{M_B} B_j^F(t) D_{ji}(R). \quad (6)$$

(c) Since the resulting representations may not be unitary, we need to compute the Hermitian

Table 2

Numbers of linearly independent elemental operators. Δ^{++} operators have uuu flavor structure, (Ω^- have sss), N^+ operators are $uud - duu$, Σ^+ are uus and suu (Ξ^0 are ssu and uss), and Λ^0 operators have $uds - sud$ flavor structure.

Operator Type	Δ, Ω	N	Σ, Ξ	Λ
single-site	20	20	40	24
singly-displaced	240	384	624	528
doubly-displaced-I	192	384	576	576
doubly-displaced-L	768	1536	2304	2304
triply-displaced-T	768	1536	2304	2304
triply-displaced-O	512	1024	1536	1536

metric matrix M

$$M_{ij} = \frac{1}{g_{O_h^D}} \sum_{R \in O_h^D} \sum_{k=1}^{M_B} D_{ki}(R)^* D_{kj}(R). \quad (7)$$

(d) For each irrep Λ , we compute the $M_B \times M_B$ projection matrix for row $\lambda = 1$:

$$P_{ij}^{\Lambda\Lambda F} = \frac{d_\Lambda}{g_{O_h^D}} \sum_{R \in O_h^D} \left[D_{\lambda\lambda}^{(\Lambda)}(R)^* D_{ji}(R) \right]_{\lambda=1}. \quad (8)$$

(e) From the rows of this projection matrix, r linearly-independent operators are obtained:

$$B_i^{\Lambda\Lambda F}(t) = \sum_{j=1}^{M_B} c_{ij}^{\Lambda\Lambda F} B_j^F(t), \quad (\lambda = 1) \quad (9)$$

$$\sum_{k,l=1}^{M_B} c_{ik}^{\Lambda\Lambda F*} M_{kl} c_{jl}^{\Lambda\Lambda F} = \delta_{ij}, \quad (i = 1 \dots r). \quad (10)$$

(f) For each of the r operators $B_i^{\Lambda\Lambda F}(t)$ in the first row $\lambda = 1$, we obtain partner operators in all other rows $\mu > 1$ using

$$c_{ik}^{\Lambda\mu F} = \sum_{j=1}^{M_B} c_{ij}^{\Lambda\Lambda F} \frac{d_\Lambda}{g_{O_h^D}} \sum_{R \in O_h^D} D_{\mu\lambda}^{(\Lambda)}(R)^* D_{kj}(R). \quad (11)$$

We average over rows for increased statistics. The total numbers of operators one obtains in each row in the different isospin channels are given in Table 3.

With such large numbers of operators, minimizing the number of quark propagator sources is crucial. We have recently finished writing software which, for each element of the baryon correlation matrices, performs all of the needed Wick contractions and simplifies the results by rotating individual terms to minimize the number of

Table 3

Total numbers of baryon operators for two different displacement lengths.

Irrep	Δ, Ω	N	Σ, Ξ	Λ
G_{1g}	221	443	664	656
G_{1u}	221	443	664	656
G_{2g}	188	376	564	556
G_{2u}	188	376	564	556
H_g	418	809	1227	1209
H_u	418	809	1227	1209

displacement orientations of the baryon sources. We hope to report on initial quenched simulations in the near future. Due to the large sizes of the baryons and their large masses, the use of improved actions on anisotropic lattices is important. Ultimately, when quark loops are included at realistically light quark masses, multi-hadron (baryon-meson) operators must be included in our correlation matrices, and finite-volume techniques will need to be employed to ferret out the baryon resonances from uninteresting scattering states. We are currently exploring different ways of building such operators.

Although we focused on tri-quark baryon operators, our approach to constructing hadronic operators is applicable to mesons, tetra-quark, and penta-quark operators. By including $\mathbf{E}$ and $\mathbf{B}$ fields into our operators, hadrons bound by an excited gluon field can also be studied. This work was supported by the U.S. National Science Foundation through grants PHY-0099450 and PHY-0300065, and by the U.S. Department of Energy under contracts DE-AC05-84ER40150 and DE-FG02-93ER-40762.

REFERENCES

1. Talk by I. Sato, these proceedings.
2. Talk by S. Basak, these proceedings.
3. LHP Collaboration, Nucl. Phys. B (Proc. Suppl.) **129**, 236 (2004).
4. C. Morningstar, M. Peardon, Phys. Rev. D **69**, 054501 (2004).